

Affordable Care Act Medicaid expansions and maternal morbidity

Pinka Chatterji¹  | Hanna Glenn² | Sara Markowitz³  | Jennifer Karas Montez⁴

¹Department of Economics, University at Albany and NBER, Albany, New York, USA

²Department of Economics, Emory University, Atlanta, Georgia, USA

³Department of Economics, Emory University and NBER, Atlanta, Georgia, USA

⁴Sociology Department, Syracuse University, Syracuse, New York, USA

Correspondence

Sara Markowitz, Department of Economics, Emory University and NBER, Atlanta, GA, USA.

Email: sara.markowitz@emory.edu

Funding information

National Institutes of Health, Grant/Award Numbers: R01 AG055481-01, R03 HD100709-02

Abstract

In this paper, we test whether the Affordable Care Act Medicaid expansions are associated with maternal morbidity. The ACA expansions may have affected maternal morbidity by increasing pre-conception access to health care, and by improving the quality of delivery care, through enhancing hospitals' financial positions. We use difference-in-difference models in conjunction with event studies. Data come from individual-level birth certificates and state-level hospital discharge data. The results show little evidence that the expansions are associated with overall maternal morbidity or indicators of specific adverse events including eclampsia, ruptured uterus, and unplanned hysterectomy. The results are consistent with prior research showing that the ACA Medicaid expansions are not statistically associated with pre-pregnancy health or maternal health during pregnancy. Our results add to this story and find little evidence of improvements in maternal health upon delivery.

KEY WORDS

Affordable Care Act, Medicaid, severe maternal morbidity

1 | INTRODUCTION

Against a backdrop of declining infant mortality rates, a discordant new trend has emerged in the United States—maternal mortality (MM) and severe maternal morbidity (SMM) rates are rising. Maternal mortality has increased by an estimated 58% over the last 2 decades, and SMM, that is, unexpected severe complications of labor and delivery, increased by 170% during the same time period (Premier, 2019; WHO, 2019). These trends are unique to the U.S., with no other industrialized countries experiencing increases in MM in recent years (WHO, 2019).

Rising rates of MM and SMM may highlight an unmet need for continuous and comprehensive health insurance coverage for women long before pregnancy begins and well after childbirth (Searing and Ross, 2019). This is particularly true for low-income older mothers, since both low SES and advanced maternal age are risk factors for adverse maternal outcomes (Dahal et al., 2017; Gray et al., 2012). Also, there is growing evidence about the importance of pre-conception health care for mothers (Barry, 2011; Johnson et al., 2006).

The primary source of insurance coverage for low-income mothers in the U.S. is Medicaid. Current federal law requires states' Medicaid programs to cover low-income pregnant women up to 138% of the federal poverty line (FPL) although many states' cutoffs exceed this minimum. As of 2018, Medicaid paid for 43% of all births (MACPAC, 2020). Prior to the Affordable Care Act (ACA) expansions, Medicaid only covered eligible pregnant women from conception to 60 days post-partum, leaving large numbers of low-income women uninsured and/or switching between insurance plans prior to conception and after the 60-day post-delivery period has ended (Daw et al., 2019).

In 2014 and later, under the ACA, 39 states (including DC) expanded Medicaid thresholds to include childless low-income adults (KFF, 2020a). Although not targeted at new mothers specifically, ACA Medicaid expansions may have improved the

continuity of Medicaid coverage from the pre-conception to the post-partum period by expanding eligibility to all adults with incomes below 138% of the FPL (Daw & Somers, 2019). This group previously had little access to Medicaid. Estimates based on pre-ACA expansion data suggest that more than half of women who had Medicaid-covered deliveries lost coverage in the months following childbirth; these women likely would retain coverage, and also likely would have had pre-conception coverage, if living in a state that expanded Medicaid under the ACA (Daw et al., 2017; KFF, 2013).

Expanding access to health insurance has the potential to increase women's access to and receipt of care prior to conception. Studies have shown that the ACA Medicaid expansions led to increases in Medicaid enrollment among women of reproductive ages, and if the new access to care results in the identification and management of chronic health conditions, women may enter pregnancy in better health (Clapp et al., 2018; Geiger et al., 2021; Johnson et al., 2018; Lyu & Wehby, 2021). In addition, there is evidence that the ACA expansions have improved hospitals' financial positions (Dranove et al., 2017; Guth et al., 2020; Moghtaderi et al., 2020). Whether the financial improvements translate to improved health outcomes is still unknown; however, one study finds that ACA expansion is associated with improvements in nurse staffing ratios and decreases in readmission rates (Tarazi, 2020).

The purpose of this paper is to test whether the ACA Medicaid expansions are associated with SMM. We envision that one causal pathway operates through women's increased enrollment in Medicaid during the pre-conception period, which in turn may affect health status during pregnancy and, ultimately, maternal health outcomes measured at the time of childbirth. Another potential pathway is Medicaid expansion improving quality of in-hospital care at delivery, leading to better outcomes at the time of childbirth. We use two sources of data on SMM: 1) individual-level data from the National Vital Statistics System Birth Certificates, and 2) state-level data on SMM rates from the Healthcare Cost and Utilization Project (HCUP). The advantage of the individual-level birth certificates data is the individual characteristics allow us to select a sample that is highly likely to be affected by the ACA Medicaid expansions. The disadvantage is the birth certificates contain only a limited set of SMM-related outcomes available for analyses. On the other hand, the state-level HCUP data provides high-quality information on SMM rates over time, but do not have information on local or individual characteristics. We use difference-in-difference models and event studies to evaluate the effects of the ACA Medicaid expansions. The results from both data sets are consistent and show little evidence that the expansions improved SMM.

2 | BACKGROUND

Severe maternal morbidity represents “unintended outcomes of the process of labor and delivery that result in significant short-term or long-term consequences to a woman's health” (Kilpatrick et al., 2016). The CDC identifies SMM through twenty-one diagnosis and procedure codes in combination with hospital discharge data to identify delivery hospitalizations (CDC, 2022). In this study, we focus on SMM rather than maternal deaths since SMM is much more common, and MM is often hard to identify in existing data sources (MacDorman et al., 2016). As we discuss below, however, our SMM measures capture some maternal deaths.

The medical causes of pregnancy-related morbidity and deaths include hemorrhage, hypertensive disorders of pregnancy, infection, embolisms, anesthesia complications, cardiovascular conditions, cardiomyopathy, cerebrovascular accident, and non-cardiovascular medical conditions (Creanga et al., 2015). What remains unknown is why SMM rates have been rising since the late 1990s (Creanga et al., 2012, 2014; Grobman et al., 2015; Moaddab et al., 2016). Much of the current research focuses on identifying individual-level, proximal risk-factors for SMM. Obesity, older maternal age, smoking, and comorbid chronic health conditions are all risk factors for SMM (Freese et al., 2020). Low socioeconomic status is also a risk factor, and racial disparities in SMM, which are well-documented, may be related to factors such as poor-quality medical care and structural racism (Grobman et al., 2014; Howell et al., 2016; Liu et al., 2019). Geography may play a role in that women residing in the most urban and the most rural areas of the U.S. also have elevated rates of SMM (Luke et al., 2021).

The proximal correlates of SMM may be useful as markers to monitor for SMM, but we do not yet have a good understanding of why many women start pregnancy in poor health or what underlying factors affect labor and delivery and lead to SMM. In a comprehensive review of maternal deaths, researchers find that the most common factors underlying preventable maternal deaths are related to patients, providers, and systems of care (Building U.S. Capacity to Review and Prevent Maternal Deaths, 2018). Patient factors include chronic conditions such as obesity, and lack of knowledge of warning signs and the need to seek care. Provider factors include delay in diagnosis and effective treatment, missed diagnosis, and ineffective treatments. System factors include inadequate training, unavailable personnel, and lack of coordination between providers in patient management.

The ACA Medicaid expansions may influence the health of the mother entering pregnancy and the quality of care during labor and delivery. If the expansions lead to more affordable and accessible health care, women may enter pregnancy in better health. If the expansions also improve hospital finances and affect the available resources, this in turn could lead to better

outcomes for women at delivery. However, it is not clear the extent to which these mechanisms could affect the relatively rare SMM outcomes, and thus we investigate the relationship empirically.

2.1 | Medicaid

Medicaid is a public health insurance program that primarily serves low-income adults, pregnant women, children, and disabled individuals. Medicaid programs are administered by states under federal guidelines and are jointly funded by states and the federal government (Medicaid, 2020). In recent years, Medicaid paid for about 43% of births in the US, with tremendous variation across states (MACPAC, 2020). For example, as of 2018–2019, Medicaid paid for 28% of births in Utah but paid for 71% of births in New Mexico (KFF, 2020b). Differences in the percentages of Medicaid-covered births across states are due to differences in income eligibility and other Medicaid program characteristics (e.g., payment rates to providers, enrollment/renewal processes) across states, as well as to differences in the SES and demographic characteristics of mothers and economic conditions across states.

Eligibility for Medicaid for pregnant women originally was tied to eligibility for Aid to Families with Dependent Children (AFDC). However, starting in the mid-1980's, changes in federal law required states to cover larger populations of low-income pregnant women and children, ultimately allowing states to cover pregnant women and children up to 185% of the federal poverty line (FPL) (Gruber, 1996). Some states expanded eligibility further without matching funds from the federal government. For low-income parents, Medicaid eligibility also expanded, but remained stringent, at less than 100% of the FPL. As a result, particularly between 1989 and 1997, there was a large increase in the percent of pregnant women eligible for Medicaid, as all states increased thresholds, although to varying degrees and at different times. Although the expansions covered prenatal care and delivery, many women were uninsured before getting pregnant and after the delivery, critical time periods for health.

The passage of the Patient Protection and ACA in 2010 led to another period of state variation in Medicaid expansions, but in this period, expansions targeted non-elderly adults up to 138% of the FPL (although the expansions extended coverage up to 133% of the FPL, this translates into 138% for most applicants since states now use Modified Adjusted Gross Income for eligibility, which involves a 5% disregard of income) (NCSL, 2011). Whereas during the prior expansions, states were expanding thresholds to varying levels, the ACA expansions involved more consistent expansions of thresholds across 39 states with most expansions occurring in 2014 or shortly after (see Table 1).

Unlike early expansions, the ACA expansions did not specifically target pregnant women, but they may still impact maternal health through expanded coverage to adults in the pre-conception period and post-delivery period. Studies have shown that the ACA Medicaid expansions led to increases in Medicaid enrollment among women of reproductive ages

TABLE 1 Affordable Care Act (ACA) Medicaid expansion dates.

Expansion in 2014
Arkansas ^a , Illinois, Iowa, Kentucky, Maryland, Michigan (April 1, 2014), Nevada, New Hampshire (August 15, 2014), New Mexico, North Dakota, Ohio, Oregon, Rhode Island ^a , Washington, West Virginia
Expansion in 2014, but with prior Medicaid for childless adults at < 138% of poverty
Arizona ^a , Colorado, Connecticut ^a , Delaware, Hawaii ^a , New Jersey ^a , New York
Expansion in 2015
Alaska (September 1, 2015), Pennsylvania, Indiana (February 1, 2015)
Expansion in 2016
Montana, Louisiana (July 1, 2016)
Expansion in 2019
Maine, Virginia
States with prior Medicaid for childless adults at ≥138% of poverty
California, District of Columbia, Massachusetts, Minnesota, Vermont

Note: Dates are as of January of the current year unless otherwise stated. States not listed are categorized as non-expansion states. States with prior Medicaid for childless adults at ≥138% of poverty are excluded from all models. States with expansion dates in the second half of the year are considered in effect for the next calendar year for the state-level SMM models.

^aStates excluded from birth certificate analyses due to lack of pre-period data resulting from late adoption of new birth certificate form: Alabama, Arizona, Arkansas, Connecticut, Hawaii, New Jersey, Rhode Island, West Virginia.

Source: Kaiser Family Foundation, Status of State Medicaid Expansion Decisions: Interactive Map. Available at: <https://www.kff.org/medicaid/issue-brief/status-of-state-medicaid-expansion-decisions-interactive-map/>. Kaiser Family Foundation, Medicaid Income Eligibility Limits for Other Non-Disabled Adults, 2011–2022. Available at: <https://www.kff.org/medicaid/state-indicator/medicaid-income-eligibility-limits-for-other-non-disabled-adults>.

(Clapp et al., 2018; Geiger et al., 2021; Johnston et al., 2018; Lyu & Wehby, 2021). If the new access to care results in the identification and management of chronic health conditions, women may enter pregnancy in better health. There is some evidence that the expansions accomplish these goals. Johnson et al. (2018) find that low-income women of reproductive age are more likely to have a personal doctor and less likely to face cost-based barriers to care after the Medicaid expansions. Myerson et al. (2020) finds that the expansion is associated with increased pre-conception health counseling and folic acid intake. However, this study along with Geiger et al. (2021), find no effects of the ACA expansions on diagnoses of diabetes or hypertension among low-income women of reproductive age.

Another pathway through which the ACA expansion may affect maternal health and SMM is via improved in-hospital financial positions and patient care. Previous research has shown that the Medicaid expansion reduced uncompensated care costs in hospitals, and that this effect was larger among hospitals that treat a disproportionate share of low-income individuals (Camilleri, 2018; Dranove et al., 2017; Guth & Ammula, 2021; Guth et al., 2020). Other studies find that the expansion is associated with improved hospital financial performance and improved operating margins, although the magnitude of the improvements may vary among urban and rural hospitals (Blavin & Ramos, 2021; Guth & Ammula, 2021; Guth et al., 2020; Moghtaderi et al., 2020; Rhodes et al., 2020). There is not much literature on how hospitals use the savings; however, Tarazi (2020) shows that the expansion is associated with improvements in nurse staffing ratios. Whether the financial improvements translate to improved health outcomes is still an open question. Tarazi (2020) finds decreases in 30-day Medicare readmission rates while Chatterjee et al. (2021) shows no improvements in hospital associated infections, mortality, and re-admission rates associated with the ACA expansion in safety-net hospitals. Nevertheless, the changes in the financial stability of hospitals and possible quality improvements may be a pathway through which the ACA expansions affect SMM.

2.2 | Previous literature on the effects of the ACA medicaid expansions on maternal health

While there exist hundreds of studies on the effects of ACA Medicaid expansions, only a few specifically focus on maternal health (Bellerose et al., 2022). Three recent papers are highly relevant to our study; studies by Palmer (2020), Margerison et al. (2021), and Eliason (2020), all of which address the question of whether the ACA Medicaid expansions affected different aspects of maternal health. Palmer (2020) and Margerison et al. (2021) both use the Vital Statistics birth certificate data to examine the effects of ACA Medicaid expansions on a variety of maternal and infant health outcomes. Margerison et al. (2021) examine measures of pre-pregnancy health (cigarette smoking, BMI, diabetes, hypertension); health and health behaviors during pregnancy (timing of prenatal care, cigarette smoking, diabetes, hypertension, eclampsia) and infant health outcomes (preterm birth and birth weight for gestational age). Palmer (2020) examines many of the same maternal and infant outcomes, along with use of anesthesia during childbirth, Cesarean section, NICU admission, and breastfeeding. Neither paper examines indicators of SMM other than eclampsia, which is studied by Margerison et al. (2021).

Palmer (2020) measures Medicaid expansion with a simulated eligibility measure reflecting the proportions of women in certain demographic groups who would be eligible under different state rules. She presents all results separately by race/ethnicity and finds that increasing the share of eligible women is associated with: (1) a small increase in first trimester prenatal care among Whites and Blacks; (2) reduced pre-pregnancy and third trimester smoking among Black women; and (3) an increase in C-sections among Black women. The Medicaid expansions are statistically unrelated to gestational diabetes and hypertension, use of anesthesia, NICU admissions, and breastfeeding.

Margerison et al. (2021) first measure Medicaid expansion by an indicator variable for conceptions that occurred at least a year post expansion versus conceptions that are timed within a year or less or no expansion. The authors also adjust the timing to reflect 6 months and 2-year windows. The second measure of expansion used is a simulated probability that the mother was Medicaid eligible prior to pregnancy. The authors' primary conclusion is that there are no improvements in maternal and infant health outcomes associated with the expansions.

By contrast, Eliason (2020) finds that the ACA expansions are associated with reductions in the rate of growth of maternal morbidity. Using data from the National Center of Health Statistics Underlying Cause of Death files and a difference-in-difference framework, Eliason estimates a reduction in maternal deaths of about 7 per 100,000 in expansion states. One limitation to this study is concerns about the accuracy of cause-of-death information on the death certificates (MacDorman et al., 2016).

Our study adds to this literature by focusing on SMM. SMM is much more common than MM and can have serious, long-term health implications for mothers and children. We recognize that the expansions can affect SMM through both pre-pregnancy health and improved quality of care at delivery. We utilize nuanced measures of Medicaid expansion by

evaluating the effects of the different length of time eligible for women without children during the pre-conception period. We present some results among different groups of women in order to see if certain women can be affected differentially by the Medicaid expansion. Overall, our results concur with Palmer (2020) and Margerison et al. (2021) and differ from those of Eliason (2020). We find no evidence that the Medicaid expansions altered SMM as defined by the incidence of SMM conditions and by state SMM rates.

3 | DATA AND MEASURES

Data on maternal health outcomes come from two sources. The first is individual-level data from the National Vital Statistics System Birth Certificates and the second is state-level data on SMM rates from Healthcare Cost and Utilization Project (HCUP). The advantage of the birth certificate data is that the individual-level data allows us to account for some individual characteristics as well as enabling us to select a sample that is likely to be affected by the ACA Medicaid expansions. The disadvantage is the limited number of health outcomes available (described below). The state-level HCUP data provides high-quality information on SMM rates over time, but without allowing us to include local or individual characteristics. The conclusions from this study will be drawn with consideration of both sets of results.

Birth Certificates: The National Vital Statistics System Birth Certificates provides individual-level data on adverse health outcomes associated with labor and delivery. Termed “SMM conditions”, these items are components of the CDC's definition of SMM (CDC, 2022). These outcomes include blood transfusion; ruptured uterus; unplanned hysterectomy; and eclampsia. We also include admission to intensive care unit, which is not part of the 21 diagnoses/procedures identified by CDC as part of SMM but nevertheless indicates that a serious adverse health event has occurred. The dependent variables we analyze are indicator variables for (1) the presence of any of these morbidities; (2) the presence of any of these morbidities excluding blood transfusion (as per an alternative definition of SMM from CDC); and (3) indicator variables for each component individually. The individual-level natality sample includes all live births including live births during which the mothers died during delivery.

We used data for births occurring in 2011–2018. These years include both pre and post periods for the ACA Medicaid expansions. We do not use earlier years of the data because earlier revisions of the birth certificate forms resulted in many states with missing values for the maternal morbidity measures as states adopted the new form. This issue also necessitates the deletion of eight states from all models due to lack of data in 2011–2013 (details are in Table 1). The estimation sample consists of twenty treatment states and eighteen control states.

In the pre-conception period, women fall into two broad Medicaid eligibility categories based on family status. If the woman has no prior births, she falls into the eligibility group of childless adults. This is the group that is specifically targeted by the ACA Medicaid expansions. Depending on the state and year, low-income women in this group may have had no options for public health insurance and may have benefited from the ACA Medicaid expansion. Women with children fall into a different Medicaid eligibility category for parents. Again, depending on the state and year, low-income women in this group may have been already eligible for Medicaid, making the expansion less relevant for them. Using information on the birth certificate data on number of prior live births, we limit the sample to women who experience a first birth in order to help identify the group of women more likely to be affected by the ACA expansions.

We also limit the sample to women ages 21 and up to avoid teenage pregnancies and the different insurance eligibility rules for these young women. Further, we limit the sample to singleton births since mothers giving birth to multiple children are likely to face different pregnancy and delivery conditions. Limiting to singletons is also useful to eliminate the potentially confounding role of assisted reproductive technology that is more commonly used by older mothers. (Figures S1–S3 in the appendix shows the main set of results for a sample limited to multiple births and they are very similar to our main findings). A difficulty to using birth certificate data is identifying women who are eligible for Medicaid. To address this, we limit the sample to women with fewer than 4 years of college, that is, those who have an associate degree, a high school degree, or less.¹ This limitation increases the likelihood that women in the sample are eligible for Medicaid, but we have no way of knowing whether those who are eligible enroll in the program. Thus, we estimate intent-to-treat effects. In auxiliary models, we further restrict the sample to unmarried women who presumably do not have spousal income and therefore may be more likely to be Medicaid eligible. Lastly, we test some models where we limit the sample to mothers ages 26 and up in order to avoid the confounding issue of the dependent coverage provision of the ACA. However, none of these sample restrictions appreciably affects the results (see Figures S4–S9 in the appendix).

The birth certificates include a limited set of maternal demographic characteristics including age, race/ethnicity, marital status, years of education. These are included in all models, and observations with missing values are preserved with

TABLE 2 Summary statistics, birth certificate data.

Variable	Mean	Std. Err.	Min	Max
Any SMM condition	0.009	0.09	0	1
Any SMM, no transfusion	0.006	0.08	0	1
Transfusion	0.003	0.06	0	1
Ruptured uterus	0.0001	0.01	0	1
Unplanned hysterectomy	0.0002	0.01	0	1
Eclampsia	0.003	0.06	0	1
Admission to ICU	0.002	0.04	0	1
ACA medicaid expansion	0.23	0.42	0	1
Months since expansion	5.48	12.10	0	55
Expansion within 1 year prior to conception	0.06	0.24	0	1
Expansion more than a year prior to conception	0.17	0.37	0	1
Age	26.11	4.58	21	50
Married	0.47	0.50	0	1
Hispanic	0.20	0.40	0	1
Black	0.19	0.39	0	1
Other race	0.06	0.23	0	1
C-section	0.33	0.47	0	1
County unemployment	6.09	2.25	0.8	20
Real income per capita	189.45	56.05	0	1002
County percent poverty	15.50	5.48	2.6	57
County bassinets per 100,000 population	18.29	14.72	0	5878
County ob/gyn and physicians per 100,000 population	82.51	34.73	0	647
Missing value on hispanic	0.004	0.06	0	1
Missing value on married	0.0001	0.01	0	1
Missing value on C-section	0.0004	0.02	0	1

Note: Data from National Vital Statistics System Birth Certificates. $N = 3,878,963$ for all variables except eclampsia ($N = 3,880,558$).

an indicator variable for missing. We also include method of delivery (C-section vs. vaginal birth) in order to account for conditions surrounding delivery (as a sensitivity check, we estimate models with and without this covariate, and also consider C-section as a dependent variable; results shown in Figures S10–S13 in the appendix and discussed below). Using the mother's county of residence available on the birth certificates, we also include time-varying controls for county-level health resources and economic conditions: the number of primary care physicians and obstetrician/gynecologists per capita, hospital capacity measured by number of bassinets per capita, the county unemployment rate, county real per capita income, and the percent of the county population living in poverty. These county-level variables all come from the Area Health Resource File.

Table 2 shows the means of the dependent variables for our sample of women with first births along with the other included variables. SMM conditions are rare, with only 0.9% of women experiencing any of the conditions. Eclampsia is the most common with 0.3% of women experiencing this condition. About 23% of the sample is observed after the ACA expansions are in effect, with 6% of the women exposed for less than a year and 17% exposed for more than a year.

State Severe Maternal Morbidity Rates: The Healthcare Cost and Utilization Project provides 'fast stats' on SMM rates by state for the years 2010–2020 (HCUP May 2022). These data are drawn from the HCUP State Inpatient Databases, which contain the universe of state hospital inpatient discharge records for patients treated in community hospitals. Rates of SMM represent the experiences of female patients ages 12–55. The numerator is the number of discharges in the state with a diagnosis or procedure indicating SMM according to the definition of CDC (2022), excluding blood transfusion. The denominator is the corresponding number of discharges indicating an in-hospital delivery. The state-level SMM rates include women who survive and who died of SMM during a hospital delivery. The rates, however, do not include cases in which the mother died during

SMM rate	N	Mean	Std. Dev.	Min	Max
All	476	69.65	14.55	33.67	122.14
Age 20–24	476	57.34	15.56	10.27	118.23
Age 25–34	476	65.41	14.29	26.00	114.59
Age 35–55	476	106.55	26.74	15.55	251.57
Medicaid and Medicare	465	77.58	17.28	25.53	134.76
Private insurance	465	63.56	15.07	22.84	113.00
Income quartile 1	475	78.31	22.27	0.00	170.75
Income quartile 2	476	70.06	18.73	28.38	185.55
Income quartile 3	476	66.59	16.17	19.41	127.52
Income quartile 4	476	63.64	29.51	0.00	467.29
Safety-net hospital	476	79.46	23.11	6.03	179.17
Non-safety-net hospital	476	63.11	15.54	13.96	114.28
Metro hospital	436	53.14	17.63	0.00	166.02
Metro teaching hospital	459	81.77	17.66	33.06	164.06
Nonmetro hospital	448	50.24	17.98	0.00	131.47
<500 deliveries	473	56.21	25.98	0.00	251.68
500-1000 deliveries	452	52.87	20.49	7.95	184.09
>1000 deliveries	454	75.26	15.81	32.16	128.43

TABLE 3 Summary statistics, state-level Healthcare Cost and Utilization Project (HCUP) data.

delivery of a cause not considered to be an SMM; we also do not capture deaths or SMM that occur during out-of-hospital deliveries.


HCUP provides SMM rates for all inpatient deliveries and by select patient and hospital characteristics. These sub-groups include age, expected payer, community income level, safety-net hospital status, hospital location/teaching status, and hospital delivery volume. Rates by race are also reported, but unfortunately not until 2016 which is after the enactment of the ACA Medicaid expansions. Three states (AK, DE, and MS) and Washington DC also have no reported SMM rates prior to 2014 and are therefore excluded. Table 3 shows the means of these SMM rates and Figures 1a and 1b shows select rates over time. The upward trends in SMM for all women and for select subgroups can be clearly seen from these figures.

ACA Medicaid Expansions: The primary variables of interest reflect Medicaid access/eligibility in the pre-conception period. We focus on the pre-conception period in order to evaluate whether insurance in this period translates to improved maternal outcomes at delivery. The mechanism we envision here is improved health going into the pregnancy. The second mechanism of improved in-hospital care via financial improvement also likely requires a time lag due to the mismatch between fiscal and calendar years, and to allow for time for hospitals adjust to the change. Using ACA expansion status at the pre-conception period would allow at least a year for hospitals to respond to the new ACA-related revenue.

When using the birth certificate data, we are able to closely match the Medicaid expansion to the preconception period by determining the conception date based on birth date and gestation weeks. We create an indicator variable equal to 1 if the mother's state of residence had expanded Medicaid to childless adults at 138% of poverty by the time of conception. We then create two indicator variables capturing time of conception relative to expansion: (1) an indicator variable for whether the statutory eligibility began within 1 year prior to conception, and (2) an indicator variable for whether statutory eligibility began more than a year prior to conception. It is unknown how long before pregnancy a woman would need to begin insurance to help improve health going into pregnancy, which is why we include both the one- and two-year lookback. Since we are able to generate an approximate conception date, we can also calculate the number of months the expansion has been in effect. We use this variable in event studies to show more specifically how time before and after the ACA expansion affects SMM.

The HCUP SMM data do not allow us to observe the birth date or conception date so the match with the ACA expansion is based off of the year of the SMM event. Since we are targeting the mother's health status going into pregnancy, we specify a one-year lag in the ACA expansion date relative to the SMM event, and alternatively a two-year lag, in order to target the correct time periods. The one- and two-year lags also allow time for adjustments to hospital finances as a result of the expansion.

FIGURE 1 (a) Average severe maternal morbidity (SMM) rates by age, over time. (b) Average SMM rates by insurance status and safety net hospital status, over time. Data from Healthcare Cost and Utilization Project (HCUP), 2010–2020. [Colour figure can be viewed at wileyonlinelibrary.com]



Information on the timing of the ACA expansion come from Kaiser Family Foundation (KFF, 2020a). The list of expansion states includes the states that did not offer Medicaid to childless adults prior to the ACA, along with the seven states that offered Medicaid but at levels smaller than 138% of poverty. There are also five states that offered Medicaid at or above 138% of poverty prior to 2014. We excluded women living in these five states from all samples. Table 1 shows the expansion status and dates for all states. We test models that exclude the states that offered Medicaid prior to 2014 but at levels smaller than 138% of poverty and states that expanded after January 2014 (see Table 1). Results are not sensitive to these changes (see Figures S14–S19 in the appendix).

4 | METHODS

We use a two-way fixed effects approach to test whether the ACA Medicaid expansions affect SMM. The estimation equation for the birth certificate data is as follows:

$$SMM_{ist} = \beta_1 ACA\ Expansion_{smc} + \beta_2 X_{ist} + \beta_3 County_{yst} + \gamma_s + \tau_c + \varepsilon_{ist} \tag{1}$$

The dependent variable (SMM_{ist}) is an indicator of SMM described above for mother i in state s and birth year t . The variable $ACA\ Expansion_{smc}$ represents one of the three forms of the Medicaid expansion variables described above based on the year of conception (c) and month of conception (m). That is, Medicaid expansion is alternatively defined by: (1) a binary variable indicating whether state had an expansion in effect in conception month and year; (2) a set of two binary variables indicating whether state had expanded less than a year ago as of conception month and year, or whether state had had expanded more than a year ago as of conception month and year (non-expander states get zeros for both indicators); and (3) number of

months expansion had been in effect as of conception month and year, including zeros for non-expanders. The model includes the vectors of maternal characteristics in birth year (X_{ist}) and county characteristics in birth year (C_{st}), state fixed effects (γ_s), and conception year fixed effects (τ_c). Including month and year of delivery fixed effects did not affect our estimates appreciably; see appendix Figures S20–S22. Excluding bassinets per capita and physician density per capita (potential mechanisms linking expansion to outcomes) also did not affect our findings in a meaningful way; see appendix Figures S23–S25. We estimate linear probability models with robust standard errors clustered on state. Logit models yield very similar results; see appendix Table S1. Since we only include 38 states in our sample, as a sensitivity check, we also estimate the standard errors using wild cluster bootstrapping with clustering on state; results are similar to the main findings and are shown in the appendix in Table S2.

The estimated coefficient(s) on the ACA expansion captures the reduced-form intent-to-treat estimate of the effect of the expansion on outcomes. Our approach controls for various economic conditions that affect insurance coverage, but we cannot specifically control for other aspects of the ACA implemented at the same time as Medicaid expansions. However, as many aspects of the ACA applied to all states, our method nets out those common trends and evaluates the effects in the states that expanded relative to the pre-expansion period.

The estimation equation for the HCUP state-level data is as follows:

$$SMM_{stg} = \beta_1 ACA\ Expansion_{st-1} + \beta_2 State\ Characteristics_{st} + \gamma_s + \tau_t + \varepsilon_{st} \quad (2)$$

Here, SMM is the state-level rate in the year of the child's birth when the SMM occurred, and the ACA expansion is an indicator variable for expansion 1 year prior to the SMM event (and alternatively 2 years prior). SMM is also analyzed among the select patient and hospital groups (g) as described above. State characteristics include the unemployment rate, real per capita income, and the percent of the state population living in poverty. These models are estimated with OLS with robust standard errors clustered on state. We also test models using the natural log of the SMM rates and results remain unchanged (see Figures S26–S28 in appendix).

The current literature on the validity of TWFE difference-in-difference models identifies three potential problems: parallel trends, control group composition, and staggered timing/treatment heterogeneity (Goodman-Bacon, 2021; Sun & Abraham, 2021). The majority of states expanded Medicaid in 2014, but there are five states that expanded in 2015/2016 and two that expanded in 2019. These later two are only within the sample time frame for the state-level HCUP data that extends to 2020.

We address these issues in a number of ways. First, we present event studies, in which the binary indicators of Medicaid expansion in Equations (1) and (2) are replaced with relative time indicators for the months leading up to the expansion, and for the months after the expansion (with the month prior to expansion left out as the baseline period). The event studies allow us to examine treatment effect dynamics as well as to test for divergent trends in outcomes between expansion and non-expansion states during the pre-treatment period (pre-trends). Second, we note that the exclusion of the states with prior Medicaid for childless adults helps to keep the control group 'clean' and comprised of only non-adopting states. We also test models that exclude the states that offered Medicaid prior to 2014 but at levels smaller than 138% of poverty. Results are not sensitive to this change (Figures S14–S16 in the appendix).

Third, we address the potential issue of bias caused by staggered timing of expansions and treatment effect heterogeneity using two different approaches: (1) by re-estimating our event study using the newly developed estimation routine by Sun and Abraham (2021); and (2) by comparing states that expanded in January 2014 to non-expanders (i.e., dropping later expanders from the sample). We compare the estimates from these two methods against our standard two-way fixed effects model and find the results are consistent across all three alternative models; these results are shown in the appendix in Figures S29–S35. In sum, these results show the standard two-way fixed effects model is robust to the heterogeneity in expansion timing in our setting. Thus, we proceed with this model to conduct the DD estimation.

5 | RESULTS

We present results from the birth certificates first. Figure 2 shows the effects of the ACA expansion by the time of conception on the probability of an SMM condition at delivery from the birth certificate data. The coefficients and confidence intervals are multiplied by 100 for ease of interpretation. Each row represents a regression with a different measure of SMM as the outcome. The results show that the ACA expansions have very small and statistically insignificant effects on most measures of SMM, indicating that the expansion is not successful in improving these maternal health outcomes at delivery. One result that stands

FIGURE 2 Effect of Affordable Care Act (ACA) expansion on probability of severe maternal morbidity (SMM). Figure shows estimated coefficient from Equation (1) on indicator of whether state had expansion in effect in conception month and year. Coefficients multiplied by 100. Figure shows 95% CI calculated using standard errors clustered on state. Models also include maternal characteristics, county characteristics, and state and conception year fixed effects (coefficients not shown). $N = 3,878,963$ ($N = 3,880,558$ for eclampsia). [Colour figure can be viewed at wileyonlinelibrary.com]





FIGURE 3 Effect of time since Affordable Care Act (ACA) expansion on probability of severe maternal morbidity (SMM). Figure shows estimated coefficients from Equation (1) on two binary variables indicating (1) whether state had expanded less than a year ago as of conception month and year or (2) whether state had had expanded more than a year ago as of conception month and year (non-expander states get zeros for both indicators). Coefficients multiplied by 100. Figure shows 95% CI calculated using standard errors clustered on state. Models also include maternal characteristics, county characteristics, and state and conception year fixed effects (coefficients not shown). $N = 3,878,963$ ($N = 3,880,558$ for eclampsia). [Colour figure can be viewed at wileyonlinelibrary.com]



out is that the coefficient on transfusion is positive and statistically significant, indicating that this procedure increases after the ACA expansion.² The magnitude is also fairly substantial at a 23% increase over the mean. This is an unexpected result that requires further investigation (discussed below). Figure 3 is similar to Figure 2 except that the indicator for ACA expansion is separated into the two time periods reflecting less than 1 year prior to conception and for one or more years prior to the conception. Again, the coefficients on most measures are very small and statistically insignificant. Figure 4 tells the same story using months eligible.

Figures 5a–g shows event studies using months as the unit of time, with thirty-six months before and forty-eight months after the policy shown (the endpoints include the accumulation of months beyond those dates). The results here confirm the more aggregated policy measures with no effects of the months since ACA expansion on the probability of any SMM (Figure 5a), any SMM excluding transfusion (Figure 5b), ruptured uterus (Figure 5c) and unplanned hysterectomy (Figure 5d). The graphs for eclampsia (Figure 5e) also shows statistically insignificant coefficients; however, there is a clear downward trend to eclampsia following the ACA expansions.




FIGURE 4 Effect of months since Affordable Care Act (ACA) expansion on probability of severe maternal morbidity (SMM). Figure shows estimated coefficient from Equation (1) on number of months expansion had been in effect as of conception month and year, including zeros for non-expanders. Coefficients multiplied by 100. Figure shows 95% CI calculated using standard errors clustered on state. Models also include maternal characteristics, county characteristics, and state and conception year fixed effects (coefficients not shown). $N = 3,878,963$ ($N = 3,880,558$ for eclampsia). [Colour figure can be viewed at wileyonlinelibrary.com]

The graph for transfusion shown in Figure 5f helps explain the positive coefficients shown in Figures 2, 3, and 4. Transfusions seem to be increasing after about 3 years post, with a positive and statistically significant coefficient showing up by the end of the period. The graph for ICU admission (Figure 5g) also shows a slight upward trend after the ACA expansion, although the coefficients are insignificant. It is not clear exactly how these increased rates of transfusions and ICU admissions might be interpreted. If taken at face value, these are negative outcomes and undesirable effects of the ACA expansions. This could occur if crowd-out of private insurance occurs and assuming private insurance generates better health outcomes than Medicaid. It is difficult to assess the health effects of the different types of insurance given the varying nature of private insurance costs and benefits prior to ACA. Also, there is currently mixed evidence on crowd-out with no consensus on whether it occurred or not (Guth et al., 2020). Conversely, if the increased rates of transfusion and ICU admissions mean that more women are getting lifesaving treatment and are less likely to die in childbirth, as shown in Eliason (2020), then these trends are not troubling. For example, transfusion is an appropriate medical response to hemorrhage, so observing more transfusions after Medicaid expansion may point to higher quality medical care upon delivery.³


As mentioned previously, we generate results from alternative models: (1) excluding states that offered Medicaid prior to 2014 but at levels smaller than 138% of poverty; (2) excluding the eight states that passed the ACA after January 2014; (3) limiting the sample to unmarried women only; (4) limiting the sample to mothers ages 26 and up. These results exhibit the same overall patterns as the main results already shown with small and statistically insignificant effects for all outcomes except for transfusion (see appendix Figures S4–S9).

In the main specifications, we control for method of delivery (C-section vs. vaginal delivery) since method of delivery may be viewed as a predetermined variable and a potential cause of SMM. On the other hand, we do not know the timing of events; a C-section could be performed after and because the mother has experienced an SMM. For this reason, in the appendix, we show results from the main models with and without the C-section covariate included (Figures S10–S12). We also show estimates of the effect of ACA Medicaid expansion on the probability of having a C-section (Figure S13). Our findings are not affected by whether or not the C-section covariate is included. The effect of Medicaid expansion on the likelihood of having a C-section is not statistically significantly different from zero. Even at the outer bounds of the 95% confidence interval, the magnitude of the effect of Medicaid expansion on C-section is small (relative to the mean). This magnitude is not meaningful from a clinical or from an economic perspective (Figure S13).


The results for the state-level SMM rates are shown in Figures 6 and 7. The first row shows the result for the SMM rate among all women, with each subsequent row containing the results for a subgroup of women. Figure 6 lags ACA expansion by 1 year and Figure 7 lags the expansion date by 2 years. The results in both models show small and statistically insignificant effects of the expansion on SMM rates among all women and among all subgroups reported. Using logged rates does not change the conclusions (see appendix Figures S26–S28). An event study is shown using SMM rates among

FIGURE 5 (a) Monthly event study, any severe maternal morbidity (SMM) indicator. (b) Monthly event study, any SMM excluding transfusion indicator. (c) Monthly event study, ruptured uterus indicator. (d) Monthly event study, unplanned hysterectomy indicator. (e) Monthly event study, Eclampsia indicator. (f) Monthly event study, transfusion indicator. (g) Monthly event study, admission to ICU indicator. All coefficients multiplied by 100. Figures show 95% CI calculated using standard errors clustered on state. Models also include maternal characteristics, county characteristics, and state and conception year fixed effects (coefficients not shown). $N = 3,878,963$; $N = 3,880,558$ for Eclampsia. [Colour figure can be viewed at wileyonlinelibrary.com]

(a)



(b)




all women in Figure 8. Again, this shows no statistically significant relationship between the ACA expansion and SMM rates in any time period.


As mentioned previously, we check for the sensitivity of the results to different classifications of expansion states. Specifically, we test models that exclude the states that offered Medicaid prior to 2014 but at levels smaller than 138% of poverty. We also run models that exclude the eight states that passed the ACA after January 2014. The results are not sensitive to these changes and still show no statistically significant effects of the ACA on the measures of SMM (see appendix Figures S14–S19).

(c)

FIGURE 5 (Continued)




(d)



6 | CONCLUSIONS

In this paper, we test whether the ACA Medicaid expansions are associated with SMM as measured by state-level rates and by specific adverse health events at delivery. The causal pathways likely operate through two channels; women's increased enrollment in Medicaid which may affect health status during pregnancy and childbirth, and improved quality of in-hospital care at delivery. We use difference-in-difference models in conjunction with event studies to evaluate the effects of the ACA Medicaid expansions. The results from both data sets are consistent and show little evidence that the expansions reduced SMM or the indicator events including eclampsia, ruptured uterus, and unplanned hysterectomy. One puzzling

FIGURE 5 (Continued)



result that arises is that Medicaid expansions are positively associated with blood transfusions. This result is not necessarily undesirable since it may simply indicate an appropriate medical response, although further research is warranted to delve into this result.

We caution that while we fail to reject the null of no effect of Medicaid expansions on all maternal outcomes other than transfusion, for some outcomes, our confidence intervals are wide, and we cannot rule out large effects in either direction. With that caution in mind, we interpret our findings in the context of closely related, recent research on this topic. Our results are consistent with Palmer (2020) and Margerison et al. (2021) who both find that the ACA Medicaid expansions are generally not statistically

(g)




FIGURE 5 (Continued)




FIGURE 6 State severe maternal morbidity (SMM) rates, Affordable Care Act (ACA) expansion lagged 1 year. Figure shows estimated coefficient from Equation (2) on one-year lagged indicator of whether state had expansion in effect in birth year (when SMM event occurred). Figure shows 95% CI calculated using standard errors clustered on state. Models also include state unemployment rate, real per capita income, percent poverty, and year and state fixed effects (coefficients not shown). The first row (labeled “all”) shows results for the overall state-level SMM rate. Each subsequent row shows results for the state-level SMM rate for that sub-group. Sample sizes shown in Table 3. [Colour figure can be viewed at wileyonlinelibrary.com]

associated with pre-pregnancy health or maternal health during pregnancy. Our results add to this story and, further, find little evidence of improvements in maternal health at the time of delivery. Most recently, however, Streenland and Wherry (2023) report that ACA Medicaid expansions in four states (vs. in four states that did not expand) are associated with reductions in post-partum hospitalizations, particularly in the first sixty days after childbirth. These effects possibly come from improved continuity of insurance coverage from childbirth into the post-partum period. These latest results suggest that future research should

FIGURE 7 State severe maternal morbidity (SMM) rates, Affordable Care Act (ACA) expansion lagged 2 year. Figure shows estimated coefficient from Equation (2) on two-year lagged indicator of whether state had expansion in effect in birth year (when SMM event occurred). Figure shows 95% CI calculated using standard errors clustered on state. Models also include state unemployment rate, real per capita income, percent poverty, and year and state fixed effects (coefficients not shown). The first row (labeled “all”) shows results for the overall state-level SMM rate. Each subsequent row shows results for the state-level SMM rate for that sub-group. Sample sizes shown in Table 3. [Colour figure can be viewed at wileyonlinelibrary.com]





FIGURE 8 Event study, state severe maternal morbidity (SMM) rate. Figure shows estimated coefficients from overall state-level SMM model on relative time indicators for the months leading up to the expansion, and for the months after the expansion (with the month prior to expansion left out as the baseline period). Figure shows 95% CI calculated using standard errors clustered on state. Models also include state unemployment rate, real per capita income, percent poverty, and state and year of birth fixed effects (coefficients not shown). $N = 476$. [Colour figure can be viewed at wileyonlinelibrary.com]



focus on the effects of ACA Medicaid expansions on maternal health more broadly, from pre-conception to pregnancy to the post-partum period, and consider both the effects of gaining insurance coverage as well as having continuous insurance coverage on maternal health.

ACKNOWLEDGMENTS

We gratefully acknowledge funding from NIH (R03 HD100709-02 and R01 AG055481-01). This research also benefited from grant, P30AG066583, Center for Aging and Policy Studies, awarded to Syracuse University, in consortium with Cornell University and the University at Albany, by the National Institute on Aging of the National Institutes of Health. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

CONFLICT OF INTEREST STATEMENT

The authors have no conflicts of interest to report.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the National Vital Statistics System and from the Healthcare Cost and Utilization Project.

ORCID

Pinka Chatterji  <https://orcid.org/0000-0001-7752-2438>

Sara Markowitz  <https://orcid.org/0000-0002-3892-2145>

ENDNOTES

- ¹ Based on CPS data, as of 2021, 96 percent of women living below 100% of the FPL, and 88 percent of women living below 200% of the FPL lacked a four-year college degree (US Census Bureau, 2023). Beginning in 2014, the birth certificates report payment source for delivery. Using this information, we calculate that of all women on Medicaid, 87% have a high school degree or less, and 94% have less than a college degree. Looking at this another way, of the women with less than a college degree, 57% have Medicaid, and of the women with a high school degree or less, 61% have Medicaid.
- ² Taking a deeper look at the data, we find that three expanding states—NY, OH and IL—had appreciably higher transfusion rates in the post vs. pre-expansion time periods, and these three states are driving the statistically significant finding in the regression.
- ³ It is also possible that a few isolated statistically significant effects (e.g., transfusions) may be spurious findings due to our consideration of so many outcomes. To address this issue, following Kling et al. (2007) and Persson and Rossin-Slater (2018), we create an SMM index. To do so, we take each individual SMM outcome and re-code each one so that a higher value represents a beneficial outcome. Then, we standardize each outcome by subtracting the mean of the outcome for the never-expander states and dividing by the never-expander state standard deviation. Next, we take an unweighted mean of the standardized SMM indicators to create the SMM index. We still find no effects of the ACA Medicaid expansions on SMM when we estimate the models using the SMM index as the dependent variable (see appendix, Table S3).

REFERENCES

- Barry, M. (2011). Preconception care at the edges of the reproductive lifespan. *Nursing for Women's Health*, 15(1), 68–74. <https://doi.org/10.1111/j.1751-486x.2011.01613.x>
- Bellerose, M., Collin, L., & Daw, J. R. (2022). The ACA medicaid expansion and perinatal insurance, health care use, and health outcomes: A systematic review: Systematic review examines the effects of expanding medicaid on insurance coverage, health care use, and health outcomes during preconception, pregnancy, and postpartum. *Health Affairs*, 41(1), 60–68. <https://doi.org/10.1377/hlthaff.2021.01150>
- Blavin, F., & Ramos, C. (2021). Medicaid expansion: Effects on hospital finances and implications for hospitals facing COVID-19 challenges: Study examines medicaid expansion effects on hospital finances and implications for hospitals facing COVID-19 challenges. *Health Affairs*, 40(1), 82–90. <https://doi.org/10.1377/hlthaff.2020.00502>
- Building U.S. (2018). Capacity to review and prevent maternal deaths. In *Report from nine maternal mortality review committees*. Retrieved from <https://www.cdcfoundation.org/sites/default/files/files/ReportfromNineMMRCs.pdf>
- Camilleri, S. (2018). The ACA Medicaid expansion, disproportionate share hospitals, and uncompensated care. *Health Services Research*, 53(3), 1562–1580. <https://doi.org/10.1111/1475-6773.12702>
- Center for Disease Control and Prevention. (2022). How does CDC identify severe maternal morbidity? Retrieved from <https://www.cdc.gov/reproductivehealth/maternalinfanthealth/smm/severe-morbidity-ICD.htm>
- Chatterjee, P., Qi, M., & Werner, R. M. (2021). Association of medicaid expansion with quality in safety-net hospitals. *JAMA Internal Medicine*, 181(5), 590–597. <https://doi.org/10.1001/jamainternmed.2020.9142>
- Clapp, M. A., James, K. E., Kaimal, A. J., & Daw, J. R. (2018). Preconception coverage before and after the affordable care act medicaid expansions. *Obstetrics and Gynecology*, 132(6), 1394–1400. <https://doi.org/10.1097/aog.0000000000002972>
- Creanga, A. A., Berg, C. J., Ko, J. Y., Farr, S. L., Tong, V. T., Bruce, F. C., & Callaghan, W. M. (2014). Maternal mortality and morbidity in the United States: Where are we now? *Journal of Women's Health*, 23(1), 3–9. <https://doi.org/10.1089/jwh.2013.4617>
- Creanga, A. A., Berg, C. J., Syverson, C., Seed, K., Bruce, F. C., & Callaghan, W. M. (2012). Race, ethnicity, and nativity differentials in pregnancy-related mortality in the United States: 1993–2006. *Obstetrics and Gynecology*, 120(2, Part 1), 261–268. <https://doi.org/10.1097/aog.0b013e31825cb87a>
- Creanga, A. A., Berg, C. J., Syverson, C., Seed, K., Bruce, F. C., & Callaghan, W. M. (2015). Pregnancy-related mortality in the United States, 2006–2010. *Obstetrics and Gynecology*, 125(1), 5–12. <https://doi.org/10.1097/01.aog.0000472713.19210.f5>
- Dahal, K., Premji, S., Patel, A., Williamson, T., Peng, M., & Metcalfe, A. (2017). Variation in maternal Co-morbidities and obstetric interventions across area-level socio-economic status: A cross-sectional study. *Paediatric & Perinatal Epidemiology*, 31(4), 274–283. <https://doi.org/10.1111/ppe.12370>
- Daw, J. R., Hatfield, L. A., Swartz, K., Sommers, B. D., & April (2017). Women in the United States experience high rates of coverage “churn” in months before and after childbirth. *Health Affairs*, 36(4), 598–606. Retrieved from <https://www.healthaffairs.org/doi/pdf/10.1377/hlthaff.2016.1241>
- Daw, J. R., Kozhimannil, K. B., & Admon, L. K. (2019). High rates of perinatal insurance churn persist after the ACA. *Health Affairs Blog*. Retrieved from <https://www.healthaffairs.org/doi/10.1377/hblog.20190913.387157/>
- Daw, J. R., & Somers, B. D. (2019). The Affordable Care Act and access to care for reproductive-aged and pregnant women in the United States, 2010–2016. *American Journal of Public Health*, 109(4), 565–571. <https://doi.org/10.2105/ajph.2018.304928>

- Dranove, D., Garthwaite, C., & Ody, C. (2017). The impact of the ACA's medicaid expansion on hospitals' uncompensated care burden and the potential effects of repeal. *Issue Brief (Public Policy Institute, American Association of Retired Persons)*, 12, 1–9. <https://www.commonwealthfund.org/publications/issue-briefs/2017/may/impact-acas-medicaid-expansion-hospitals-uncompensated-care>
- Eliason, E. L. (2020). Adoption of Medicaid expansion is associated with lower maternal mortality. *Women's Health Issues*, 30(3), 147–152. <https://doi.org/10.1016/j.whi.2020.01.005>
- Freese, K. E., Bodnar, L. M., Brooks, M. M., McTigue, K., & Himes, K. P. (2020). Population-attributable fraction of risk factors for severe maternal morbidity. *American Journal of Obstetrics and Gynecology MFM*, 2(1), 100066. <https://doi.org/10.1016/j.ajogmf.2019.100066>
- Geiger, C. K., Sommers, B. D., Hawkins, S. S., & Cohen, J. L. (2021). Medicaid expansions, preconception insurance, and unintended pregnancy among new parents. *Health Services Research*, 56(4), 691–701. <https://doi.org/10.1111/1475-6773.13662>
- Goodman-Bacon, A. (2021). Difference-in-differences with variation in treatment timing. *Journal of Econometrics*, 225(2), 254–277. <https://doi.org/10.1016/j.jeconom.2021.03.014>
- Gray, K. E., Wallace, E. R., Nelson, K. R., Reed, S. D., & Schiff, M. A. (2012). Population-based study of risk factors for severe maternal morbidity. *Paediatric & Perinatal Epidemiology*, 26(6), 506–514. <https://doi.org/10.1111/ppe.12011>
- Grobman, W. A., Bailit, J. L., Rice, M. M., Wapner, R. J., Reddy, U. M., Varner, M. W., Thorp, J. M., Leveno, K. J., Caritis, S. N., Iams, J. D., Saade, G., Rouse, D. J., Blackwell, S. C., Tolosa, J. E., VanDorsten, J. P., & Tita, A. T. (2015). Racial and ethnic disparities in maternal morbidity and obstetric care. *Obstetrics & Gynecology*, 125(6), 1460–1467. <https://doi.org/10.1097/aog.0000000000000735>
- Grobman, W. A., Bailit, J. L., Rice, M. M., Wapner, R. J., Varner, M. W., Thorp, J. M., Saade, G., Caritis, S. N., Iams, J. D., Tita, A. T., Sorokin, Y., Rouse, D. J., Tolosa, J. E., & Van Dorsten, J. P. (2014). Can differences in obstetric outcomes be explained by differences in the care provided? The MFMU network APEX study. *American Journal of Obstetrics and Gynecology*, 211(2), 147.e1–147.e16. <https://doi.org/10.1016/j.ajog.2014.03.017>
- Gruber, J. (1996). *Health insurance for poor women and children in the US: Lessons from the past decade*. National Bureau of Economic Research Working Paper.
- Guth, M., & Ammula, M. (2021). Building on the evidence base: Studies on the effects of medicaid expansion, February 2020 to March 2021. In *Kaiser family foundation*.188
- Guth, M., Garfield, R., Rudowitz, R., & March (2020). The effects of medicaid expansion under the ACA: Updated findings from a literature review. *Henry Journal Kaiser Family Foundation*, 37(6), 944–950.
- HCUP Fast Stats. (2022). *Healthcare cost and utilization Project (HCUP)*. Agency for Healthcare Research and Quality. Retrieved from www.hcup-us.ahrq.gov/faststats/smm/smmquery.jsp?radio2=on&location1=US&characteristic1=01C11&location2=&characteristic2=01C11&expansionInfoState=hide&dataTablesState=hide&definitionsState=hide&exportState=hide
- Howell, E. A., Egorova, N., Balbierz, A., Zeitlin, J., & Hebert, P. L. (2016). Black-White differences in severe maternal morbidity and site of care. *American Journal of Obstetrics and Gynecology*, 214(1), 122.e1–122.e7. <https://doi.org/10.1016/j.ajog.2015.08.019>
- Johnson, K., Posner, S. F., Biermann, J., Cordero, J. F., Atrash, H. K., Parker, C. S., Boulet, S., & Curtis, M. G. (2006). Recommendations to improve pre-conception health and health care: A report of the CDC/ATSDR preconception care workgroup and the select panel on preconception care. *Morbidity and Mortality Weekly Report*, 55 RR06, 1–23.
- Johnston, E. M., Strahan, A. E., Joski, P., Dunlop, A. L., & Adams, E. K. (2018). Impacts of the affordable care act's medicaid expansion on women of reproductive age: Differences by parental status and state policies. *Women's Health Issues*, 28(2), 122–129. <https://doi.org/10.1016/j.whi.2017.11.005>
- Kaiser Family Foundation (KFF). (2013). Health care reform: Implication for women's access to coverage and care. Retrieved from https://www.kff.org/wp-content/uploads/2012/03/7987-03-health-reform-implications-for-women_s-access-to-coverage-and-care.pdf
- Kaiser Family Foundation (KFF). (2020b). Births financed by Medicaid. Retrieved from <https://www.kff.org/medicaid/state-indicator/births-financed-by-medicaid/?currentTimeframe=0&sortModel=%7B%22collId%22:%22Location%22,%22sort%22:%22asc%22%7D>
- Kaiser Family Foundation (KFF). (2020a). Status of state Medicaid expansion decisions Available at. Retrieved from <https://www.kff.org/medicaid/issue-brief/status-of-state-medicaid-expansion-decisions-interactive-map/>
- Kilpatrick, S. K., Ecker, J. L., American College of Obstetricians, & Gynecologists (2016). Severe maternal morbidity: Screening and review. *American Journal of Obstetrics and Gynecology*, 215(3), B17–B22. <https://doi.org/10.1016/j.ajog.2016.07.050>
- Kling, J. R., Leibman, J. B., & Katz, L. F. (2007). Experimental analysis of neighborhood effects. *Econometrica*, 75(1), 83–119. <https://doi.org/10.1111/j.1468-0262.2007.00733.x>
- Liu, S. Y., Fiorentini, C., Bailey, Z., Huynh, M., McVeigh, K., & Kaplan, D. (2019). Structural racism and severe maternal morbidity in New York State. *Clinical Medicine Insights: Women's Health*, 12, 1179562X19854778. <https://doi.org/10.1177/1179562x19854778>
- Luke, A. A., Huang, K., Lindley, K. J., Carter, E. B., & Joynt Maddox, K. E. (2021). Severe maternal morbidity, race, and rurality: Trends using the national inpatient sample, 2012–2017. *Journal of Women's Health*, 30(6), 837–847. <https://doi.org/10.1089/jwh.2020.8606>
- Lyu, W., & Wehby, G. L. (2021). Heterogeneous effects of affordable care act medicaid expansions among women with dependent children by state-level pre-expansion eligibility. *Journal of Women's Health*, 30(9), 1278–1287. <https://doi.org/10.1089/jwh.2020.8776>
- MacDorman, M. F., Declercq, E., Cabral, H., & Morton, C. (2016). Recent increases in the US maternal mortality rate: Disentangling trends from measurement issues. *Obstetrics & Gynecology*, 128(3), 447–455. <https://doi.org/10.1097/aog.0000000000001556>
- MACPAC. (2020). Medicaid's role in financing maternity care. Retrieved from <https://www.macpac.gov/wp-content/uploads/2020/01/Medic-aid%E2%80%99s-Role-in-Financing-Maternity-Care.pdf>
- Margerison, C. E., Kaestner, R., Chen, J., & MacCallum-Bridges, C. (2021). Impacts of Medicaid expansion before conception on pre-pregnancy health, pregnancy health, and outcomes. *American Journal of Epidemiology*, 190(8), 1488–1498. <https://doi.org/10.1093/aje/kwaa289>
- Medicaid. (2020). Retrieved from <https://www.medicaid.gov/medicaid/index.html>

- Moaddab, A., Dildy, G. A., Brown, H. L., Bateni, Z. H., Belfort, M. A., Sangi-Haghpeykar, H., & Clark, S. L. (2016). Health care disparity and state-specific pregnancy-related mortality in the United States, 2005–2014. *Obstetrics and Gynecology*, *128*(4), 869–875. <https://doi.org/10.1097/aog.0000000000001628>
- Moghtaderi, A., Pines, J., Zocchi, M., & Black, B. (2020). The effect of Affordable Care Act Medicaid expansion on hospital revenue. *Health Economics*, *29*(12), 1682–1704. <https://doi.org/10.1002/hec.4157>
- Myerson, R., Crawford, S., & Wherry, L. R. (2020). Medicaid expansion increased preconception health counseling, folic acid intake, and postpartum contraception: Study examines the impact of ACA Medicaid expansion on health behaviors including birth control use and pregnancy intention, and receipt of preconception health services. *Health Affairs*, *39*(11), 1883–1890. <https://doi.org/10.1377/hlthaff.2020.00106>
- National Conference of State Legislatures (NCSL). (2011). Medicaid and the affordable care Act.
- Palmer, M. (2020). Preconception subsidized insurance: Prenatal care and birth outcomes by race/ethnicity. *Health Economics*, *29*(9), 1013–1030. <https://doi.org/10.1002/hec.4116>
- Persson, P., & Rossin-Slater, M. (2018). Family ruptures, stress, and the mental health of the next generation. *The American Economic Review*, *108*(4–5), 1214–1252. <https://doi.org/10.1257/aer.20141406>
- Premier Inc. (2019). Bundle of joy: Maternal and infant health trends. Downloaded 10/11/2021 Retrieved from <https://explore.premierinc.com/MaternalHealthTrends>
- Rhodes, J. H., Buchmueller, T. C., Levy, H. G., & Nikpay, S. S. (2020). Heterogeneous effects of the ACA Medicaid expansion on hospital financial outcomes. *Contemporary Economic Policy*, *38*(1), 81–93. <https://doi.org/10.1111/coep.12428>
- Searing, A., Ross, D. C., & May (2019). *Medicaid expansion fills gaps in maternal health coverage leading to healthier mothers and babies*. Georgetown University Health Policy Institute, Center for Children and Families.
- Streenland, M. W., & Wherry, L. R. (2023). Medicaid expansion led to reductions in postpartum hospitalizations. *Health Affairs*, *42*(1), 18–25. <https://doi.org/10.1377/hlthaff.2022.00819>
- Sun, L., & Abraham, S. (2021). Estimating dynamic treatment effects in event studies with heterogenous treatment effects. *Journal of Econometrics*, *225*(2), 175–199. <https://doi.org/10.1016/j.jeconom.2020.09.006>
- Tarazi, W. W. (2020). Associations between Medicaid expansion and nurse staffing ratios and hospital readmissions. *Health Services Research*, *55*(3), 375–382. <https://doi.org/10.1111/1475-6773.13273>
- U.S. Census Bureau, Current Population Survey, 2023 Annual Social and Economic Supplement (CPS ASEC). Retrieved from <https://www.census.gov/data/tables/time-series/demo/income-poverty/cps-pov/pov-09.html>
- WHO. (2019). *Trends in maternal mortality 2000 to 2017: Estimates by WHO, UNICEF, UNFPA, world bank group and the united nations population division*. World Health Organization. Licence: CC BY-NC-SA 3.0 IGO.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Chatterji, P., Glenn, H., Markowitz, S., & Montez, J. K. (2023). Affordable Care Act Medicaid expansions and maternal morbidity. *Health Economics*, *32*(10), 2334–2352. <https://doi.org/10.1002/hec.4724>